

Environmental preprocessor

This is the approved revision of this page; it is not the most recent. View the most recent revision.

The printable version is no longer supported and may have rendering errors. Please update your browser bookmarks and please use the default browser print function instead.

Sources

FAO, WRI, WDI, Global_Footprint_Network,

Environmental series pulled into the model

DataDict

Table	Source	Last IFs Update	UsedInPreprocessorFileName
SeriesEnvAvgAnnTemp	http://www.cgd.ucar.edu/cas/wigley/magicc/ and http://na.unep.net/globalpop/1-degree/description.php	2009/07/27	ENV
SeriesEnvAvgAnnTempChg	http://www.cgd.ucar.edu/cas/wigley/magicc/ and http://na.unep.net/globalpop/1-degree/description.php	2009/07/27	ENV
SeriesEnvPrecipitation	http://www.cgd.ucar.edu/cas/wigley/magicc/ and http://na.unep.net/globalpop/1-degree/description.php	2009/07/27	ENV
SeriesEnvPrecipitationChg	http://www.cgd.ucar.edu/cas/wigley/magicc/ and http://na.unep.net/globalpop/1-degree/description.php	2009/07/27	ENV
SeriesEnvPM10	WDI 2014 May BATCH PULL	2014/06/11	ENV
SeriesEmissionsCarbonCDIAC	Carbon Dioxide Information Analysis Center	2013/08/06	ENV
SeriesEnvSolidFuelsImproved	Randall Kuhn prepared it from GACC tables and remove the ones that were implausible and for tiny countries. Then added regional figures based on the old WHO data, basically taking the lower of the rural and urban figures.	2013/12/19	ENV

Retrieved from "https://pardeewiki.du.edu//index.php?title=Environmental_preprocessor&oldid=1184"

This page was last edited on 2 August 2016, at 21:13.